




Inhibiteurs calciques




Les médicaments agissant sur la guanylate cyclase








sGC : Une cible, deux classes de molécules






Diurétique

Acetazolamide



Acetazolamide



Acetazolamide

- Paresthésies : 30-90 %
- Polyurie 8-55 %
- Nausées 20 %
 - > intérêt d'un test préalable

Low et al., BMJ 2012
Ellsworth et al, Am J Med 1987

Hypnotiques



In the Zzz zone: effect of z-drugs

DCI	Présentation	Doses	Tmax	½ vie	Tolérance, dépendance	Effets sur les fonctions cognitives
Zolpidem	Stilnox [®] , génériques	5-10 mg	1-2 h	2 h	rare	+/-, dose dépendants Hallucinations, amnésie, parasomnie
Zopiclone	Imovane [®] , génériques	3,75-7,5 mg	1,5-2 h	5-6 h	rare	+, dose dépendants

FDA Drug Safety Communication: Risk of next-morning impairment after use of insomnia drugs; FDA requires lower recommended doses for certain drugs containing zolpidem [1-10-2013]



Zolpidem Dosing Recommendations for Adults (Non-Elderly)

	Dosing recommendations in current drug label for zolpidem	FDA's proposed new dosing recommendations for zolpidem
Ambien, Edluar, Zolpimist	Men and Women: 10 mg once daily, immediately before bedtime	Women: 5 mg once daily, immediately before bedtime Men: 5 or 10 mg once daily, immediately before bedtime



